

THE UNITED STRATES OF AMERICA

To all to whom these presents shall come: Texas Agricultural Experiment Station

A LCCCAS, THERE HAS BEEN PRESENTED TO THE

Secretary of Agriculture

AN APPLICATION REQUESTING A CERTIFICATE OF PROTECTION FOR AN ALLEGED DISTINCT VARIETY OF SEXUALLY REPRODUCED, OR TUBER PROPAGATED PLANT, THE NAME AND DESCRIPTION OF WHICH ARE CONTAINED IN THE APPLICATION AND EXHIBITS, A COPY OF WHICH IS HEREUNTO ANNEXED AND MADE A PART HEREOF, AND THE VARIOUS REQUIREMENTS OF LAW IN SUCH CASES MADE AND PROVIDED HAVE BEEN COMPLIED WITH, AND THE TITLE THERETO IS FROM THE RECORDS OF THE PLANT VARIETY PROTECTION OFFICE, IN THE APPLICANT(S) INDICATED IN THE SAID COPY, AND WHEREAS, UPON DUE EXAMINATION MADE, THE SAID APPLICANT(S) IS (ARE) ADJUDGED TO BE ENTITLED TO A CERTIFICATE OF PLANT VARIETY PROTECTION UNDER THE LAW.

NOW, THEREFORE, THIS CERTIFICATE OF PLANT VARIETY PROTECTION IS TO GRANT UNTO THE SAID APPLICANT(S) AND THE SUCCESSORS, HEIRS OR ASSIGNS OF THE SAID APPLICANT(S) FOR THE TERM OF TWENTY YEARS FROM THE DATE OF THIS GRANT, SUBJECT TO THE PAYMENT OF THE REQUIRED FEES AND PERIODIC REPLENISHMENT OF VIABLE BASIC SEED OF THE VARIETY IN A PUBLIC REPOSITORY AS PROVIDED BY LAW, THE RIGHT TO EXCLUDE OTHERS FROM SELLING THE VARIETY, OR OFFERING IT FOR SALE, OR REPRODUCING IT, OR CORTING IT, OR EXPORTING IT, OR CONDITIONING IT FOR PROPAGATION, OR STOCKING IT FOR ANY OF THE VE PURPOSES, OR USING IT IN PRODUCING A HYBRID OR DIFFERENT VARIETY THEREFROM, TO THE EXTENT DED BY THE PLANT VARIETY PROTECTION ACT. IN THE UNITED STATES SEED OF THIS VARIETY BE SOLD BY VARIETY NAME ONLY AS A CLASS OF CERTIFIED SEED AND (2) SHALL CONFORM TO THE ACCURACY.

WHEAT, COMMON

'TAM 112'

In Testimony Thereof, I have hereunto set my hand and caused the seal of the Hant Inricty Frotestion Office to be affixed at the City of Washington, D.C. this sixth day of February, in the year two thousand and seven.

Attest:

Gamissioner Commissioner

Commissioner Plant Variety Protection Office Agricultural Marketing Service Secretary of Agriculture

U.S. DEPARTMENT OF AGRICULTURE AGRICULTURAL MARKETING SERVICE SCIENCE AND TECHNOLOGY - PLANT VARIETY PROTECTION OFFICE APPLICATION FOR PLANT VARIETY PROTECTION CERTIFICATE

The following statements are made in accordance with the Privacy Act of 1974 (5 U.S.C. 552a) and the Paperwork Reduction Act (PRA) of 1995.

SCIENCE AND TECHNOLOGY - F	LANT VARIETY P	ROTECTION OFFICE	Application in required in order to deter-	mina if a	plant unright, protection cortificate in to be included
APPLICATION FOR PLANT VA (Instructions and information co					olant variety protection certificate is to be issued until certificate is issued (7 U.S.C. 2426).
1. NAME OF OWNER			2. TEMPORARY DESIGNATION OR	3. VA	RIETY NAME
Texas Agricultural Expe	riment St	cation	TX98V9628	TAI	M 112
4. ADDRESS (Street and No., or R.F.D. No., City,	State, and ZIP Co.	de, and Country)	5. TELEPHONE (include area code)		FOR OFFICIAL USE ONLY
Office of the Director			979-845-4747 6. FAX (include area code)	PVPO	NUMBER 10600274
2147 TAMU, College Stat	ion, TX	77843-2147	979-458-4765	FILING	G DATE -
 IF THE OWNER NAMED IS NOT A "PERSON", ORGANIZATION (corporation, partnership, asso 		8. IF INCORPORATED, GIVE STATE OF INCORPORATION	9. DATE OF INCORPORATION	1	August 22, 2006
Texas State Research Ac	jency				
10. NAME AND ADDRESS OF OWNER REPRESE	ENTATIVE(S) TO S	BERVE IN THIS APPLICATION. (First p	erson listed will receive all papers)	E E S	FILING AND EXAMINATION FEES:
Janie Hurley, Licensing	Mgr., Of	fice of Tech. Com	mercialization	R E	DATE 8-22-2006 CERTIFICATION FEE:
The Texas A&M University	y System,	3369 TAMU		C E I	s 768. ∞
College Station, TX 7784	13-3369			V E D	DATE 12-12- 2006
11. TELEPHONE (Include area code)	12. FAX (Includ	e area code)	13. E-MAIL		
979-847-8682 m	1. 2/2 04	5-1402	jhurley@tam	u.eć	lu
14. CROPKIND (Common Name) wheat	16. FAMILY NA	•	18. DOES THE VARIETY CONTA	IN ANY T	RANSGENES? (OPTIONAL)
wileat	Poacea	e	☐ YES 🛱 NO		
15. GENUS AND SPECIES NAME OF CROP	17. IS THE VAR	RIETY A FIRST GENERATION HYBRID	IF SO, PLEASE GIVE THE AS	SSIGNED EREGUI) USDA-APHIS REFERENCE NUMBER FOR THE LATE THE GENETICALLY MODIFIED PLANT FOR
Triticum aestivum L.	☐ YES	Ď X no	COMMERICALIZATION.		
19. CHECK APPROPRIATE BOX FOR EACH ATTA (Follow instructions on reverse)	L ACHMENT SUBMI	ITED			EED OF THIS VARIETY BE SOLD AS A CLASS 83(a) of the Plant Variety Protection Act)
a. Exhibit A. Origin and Breeding History	of the Variety				and 22 below) 🔲 NO (If "no", go to item 23)
b. Exhibit B. Statement of Distinctness			21. DOES THE OWNER SPECIFY NUMBER OF CLASSES?	THAT S	EED OF THIS VARIETY BE LIMITED AS TO
c. 🕅 Exhibit C. Objective Description of Var	iety		X YES □ NO		
d. 💹 Exhibit D. Additional Description of the				™ =011	NDATION KETREGISTERED KETTIFIED
e. Exhibit E. Statement of the Basis of the		nip		THAT S	EED OF THIS VARIETY BE LIMITED AS TO
f. 🛱 Exhibit F. Declaration Regarding Depo	sit		X YES □ NO		
g. Xi Voucher Sample (3,000 viable untreate that tissue culture will be deposited and	ed seeds or, for tub I maintained in an a	er propagated varieties, verification approved public repository)	IF YES, SPECIFY THE NUMBI	ER 1,2,3,	etc. FOR EACH CLASS.
g. X Filing and Examination Fee (\$4,382), m States" (Mail to the Plant Variety Protect		easurer of the United	3 FOUNDATION 5 REG		ED D CERTIFIED lease use the space indicated on the reverse.)
23. HAS THE VARIETY (INCLUDING ANY HARVES FROM THIS VARIETY BEEN SOLD, DISPOSED OTHER COUNTRIES?	TED MATERIAL) O OF, TRANSFERE	OR A HYBRID PRODUCED RED, OR USED IN THE U. S. OR	24. IS THE VARIETY OR ANY CO	MPONE	NT OF THE VARIETY PROTECTED BY PLANT BREEDER'S RIGHT OR PATENT)?
□ YES □ NO			□ YES X) NO		
IF YES, YOU MUST PROVIDE THE DATE OF F FOR EACH COUNTRY AND THE CIRCUMSTA			IF YES, PLEASE GIVE COUNT REFERENCE NUMBER. (Pleas		E OF FILING OR ISSUANCE AND ASSIGNED pace indicated on reverse.)
The owners declare that a viable sample of basifor a tuber propagated variety a tissue culture w The undersioned owner(s) is(are) the owner of it	ill be deposited in	a public repository and maintained for	he duration of the certificate.		with such regulations as may be applicable, or
entitled to protection under the provisions of Section	ion 42 of the Plant	Variety Protection Act.	and Delivering and the valuety is new, the	mos, uciti	son, and occord to required in occurrate, with is

Owner(s) is (are) informed that false representation herein can jeopardize protection and result in penalties.

Maha Jusse		SIGNATURE OF OWNER		
Mark A. Hussey		NAME (Please print or type)		
CAPACITY OR TITLE	DATE	CAPACITY OR TITLE	DATE	
Associate Director, TAES	8-10-2006		tu.	

See reverse for instructions and information collection burden statement,

200600274

GENERAL INSTRUCTIONS: To be effectively filed with the Plant Variety Protection Office (PVPO), ALL of the following items must be received in the PVPO: (1) Completed application form signed by the owner; (2) completed exhibits A, B, C, E, F; (3) for a tuber reproduced variety, verification that a viable (in the sense that it will reproduce an entire plant) tissue culture will be deposited and maintained in an approved public repository; and (4) payment by credit card or check drawn on a U.S. bank for \$4,382 (\$518 filing fee and \$3,864 examination fee), payable to "Treasurer of the United States" (See Section 97.6 of the Regulations and Rules of Practice). NEW: With the application for a seed reproduced variety or by direct deposit soon after filing, the applicant must provide at least 3,000 viable untreated seeds of the variety per se, and for a hybrid variety at least 3,000 untreated seeds of each line necessary to reproduce the variety. Partial applications will be held in the PVPO for not more than 90 days; then returned to the applicant as un-filed. Mail application and other requirements to Plant Variety Protection Office, AMS, USDA, Room 401, NAL Building, 10301 Baltimore Avenue, Beltsville, MD 20705-2351. Retain one copy for your files. All items on the face of the application are self explanatory unless noted below. Corrections on the application form and exhibits must be initiated and dated. DO NOT use masking materials to make corrections. If a certificate is allowed, you will be requested to send a payment by credit card or check payable to "Treasurer of the United States" in the amount of \$768 for issuance of the certificate. Certificates will be issued to owner, not licensee or agent.

NOTES: It is the responsibility of the applicant/owner to keep the PVPO informed of any changes of address or change of ownership or assignment or owner's representative during the life of the application/certificate. The fees for filing a change of address; owner's representative; ownership or assignment; or any modification of owner's name is specified in Section 97.175 of the regulations. (See Section 101 of the Act, and Sections 97.130, 97.131, 97.175(h) of the Regulations and Rules of Practice.)

Plant Variety Protection Office

Telephone: (301) 504-5518 FAX: (301) 504-5291

General E-mail: PVPOmail@usda.gov

Homepage: http://www.ams.usda.gov/science/pvpo/PVPindex.htm

SPECIFIC INSTRUCTIONS:

To avoid conflict with other variety names in use, the applicant must check the appropriate recognized authority and **provide evidence** that the permanent name of the application variety (even if it is a parental, inbred line) has been cleared by the appropriate recognized authority before the Certificate of Protection is issued. For example, for agricultural and vegetable crops, contact: U.S. Department of Agriculture, Agricultural Marketing Service, Livestock and Seed Programs, **Seed Regulatory and Testing Branch**, 801 Summit Crossing Place, Suite C, Gastonia, North Carolina 28054-2193 Telephone: (704) 810-8870. http://www.ams.usda.gov/lsg/seed.htm.

ITEM

19a. Give:

- (1) the genealogy, including public and commercial varieties, lines, or clones used, and the breeding method;
- (2) the details of subsequent stages of selection and multiplication;
- (3) evidence of uniformity and stability; and
- (4) the type and frequency of variants during reproduction and multiplication and state how these variants may be identified
- 19b. Give a summary of the variety's distinctness. Clearly state how this application variety may be distinguished from all other varieties in the same crop. If the new variety is most similar to one variety or a group of related varieties:
 - (1) identify these varieties and state all differences objectively;
 - (2) attach replicated statistical data for characters expressed numerically and demonstrate that these are clear differences; and
 - (3) submit, if helpful, seed and plant specimens or photographs (prints) of seed and plant comparisons which clearly indicate distinctness.

19c. Exhibit C forms are available from the PVPO Office for most crops; specify crop kind. Fill in Exhibit C (Objective Description of Variety) form as completely as possible to describe your variety.

- 19d. Optional additional characteristics and/or photographs. Describe any additional characteristics that cannot be accurately conveyed in Exhibit C. Use comparative varieties as is necessary to reveal more accurately the characteristics that are difficult to describe, such as plant habit, plant color, disease resistance, etc.
- 19e. Section 52(5) of the Act requires applicants to furnish a statement of the basis of the applicant's ownership. An Exhibit E form is available from the PVPO.
- 20. If "Yes" is specified (seed of this variety be sold by variety name only, as a class of certified seed), the applicant MAY NOT reverse this affirmative decision after the variety has been sold and so labeled, the decision published, or the certificate issued. However, if "No" has been specified, the applicant may change the choice. (See Regulations and Rules of Practice, Section 97.103).
- 23. See Sections 41, 42, and 43 of the Act and Section 97.5 of the regulations for eligibility requirements.
- 24. See Section 55 of the Act for instructions on claiming the benefit of an earlier filing date.
- 22. CONTINUED FROM FRONT (Please provide a statement as to the limitation and sequence of generations that may be certified.)

23. CONTINUED FROM FRONT (Please provide the date of first sale, disposition, transfer, or use for each country and the circumstances, if the variety (including any harvested material) or a hybrid produced from this variety has been sold, disposed of, transferred, or used in the U.S. or other countries.)

August 31, 2005, Foundation seed of TAM 112 was transferred to Watley Seed Company for planting and production under a license agreement between Watley and TAES/A&M System.(U.S.)

24. CONTINUED FROM FRONT (Please give the country, date of filing or issuance, and assigned reference number, if the variety or any component of the variety is protected by intellectual property right (Plant Breeder's Right or Patent).)

According to the Paperwork Reduction Act of 1995, an agency may not conduct or sponsor, and a person is not required to respond to a collection of information unless it displays a valid OMB control number. The valid OMB control number for this information collection is 0581-0055. The time required to complete this information collection is estimated to average 1.4 hours per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information.

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, age, disability, and where applicable, sex, marital status, familial status, parental status, religion, sexual orientation, genetic information, political beliefs, reprisal, or because all or part of an individual's income is derived from any public assistance program (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at (202) 720-2600 (voice and TDD).

To file a complaint of discrimination, write to USDA, Director, Office of Civil Rights, 1400 Independence Avenue, S.W., Washington, D.C. 20250-9410, or cell (800) 795-3272 (voice) or (202) 720-6382 (TDD). USDA is an equal opportunity provider and employer.

Origin and Breeding History - 'TAM 112' Wheat

'TAM 112' (experimental designation TX98V9628) is a hard red winter wheat (Triticum aestivum L.) experimental line developed by the Texas Agricultural Experiment Station (TAES). 'TAM 112' is an F4 derived line from the cross U1254-7-9-2-1/TXGH10440. David Worrall, TAES Vernon, made the cross during the winter of 1991-92. U1254-7-9-2-1 is a USDA-ARS germplasm line from the Plant Science and Entomology Research unit, Manhattan, Kansas. It was developed from the cross TAM 200/TA2460. TA2460 is an Aegilops tauschii (one of the progenitor species of modern hexaploid wheat) line that was identified as having a unique gene for leaf rust resistance (this gene has since been designated lr41). TXGH10440 has the same pedigree as TAM 110 (TAM 105*4/Amigo*4//Largo). Like TAM 110, TXGH10440 carries gb2 gene for greenbug resistance from the germplasm line Amigo and gb3 from the synthetic hexaploid line named Largo. The F₃ and F₄ generations were grown as bulk populations on the TAES farm at Chillicothe in 1995 (year of harvest) and 1996, respectively. Random heads were harvested from the F₄ population and were planted as head-rows at Chillicothe in the fall of 1996. The line that became TX98V9628 ('TAM 112') was visually selected for its agronomic characteristics and was grown as a single plot in 1998 and in replicated trials thereafter. The selection criteria were forage potential and grain yield. It was tested in Texas Elite trials (NTE, WTE, TXE) in 2002 and 2003, Southern Regional Performance Nursery (SRPN) in 2002 and 2003, and the Texas Uniform Variety Trial (UVT) in 2004. A head-row purification was conducted in 2003. One hundred-fifty rows were visually evaluated for uniformity and 135 were harvested individually. After harvest, 10 seed each from these 135 lines were evaluated for resistance to biotype E greenbug. Based on purity for resistance, the seed from 133 of these lines were bulked. Three acres of the purified TX98V9628 was planted by Texas Foundation Seed Service of TAES in the fall of 2003 to produce Breeders seed.

TAM 112 has been observed for 6 generations during testing and seed increase, and is stable and uniform. A tall variant (10 cm taller) was identified in the breeders seed of TAM 112 at a frequency of approximately 0.01%, but was identical to the variety in all other characteristics. Up to 0.05% variant plants may be encountered in subsequent generations. This variant is commercially acceptable and predictable.

Statement of Distinctness - 'TAM 112' Wheat

'TAM 112' (TX98V9628) is awned, red-glumed, and has good potential for both forage and grain production in the southern Great Plains. It is medium height, medium-early maturing, and has winter hardiness similar to most commonly grown hard winter wheat cultivars. It is particularly suited for dryland production systems in the High Plains of Texas and similar areas in adjacent states. It is suitable for both dual purpose (grazing plus grain) and grain only systems.

TAM 112 is most similar to TAM 110. Specifically, TAM 112 and TAM 110 are similar in area of adaptation, drought tolerance, and disease and insect resistance; however, TAM 112 differs from TAM 110 in the following physical characteristics: TAM 112 has purple anthers where TAM 110 has yellow anthers; and coleoptile anthocyanin is present in TAM 112, but is absent in TAM 110. These differences are recorded in the objective descriptions for each variety.

REPRODUCE LOCALLY. Include form number and date on all reproductions. Form Approved OMB NO

According to the Paperwork Reduction Act of 1995, an agency may not conduct or sponsor, and a person is not required to respond to a collection of information unless it displays a valid OMB control number. The valid OMB control number for this information collection is 0581-0055. The time required to complete this information collection is estimated to average 2.5 hours per response, including the time for

searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, gender, religion, age, disability, sexual orientation, marital or family status, political beliefs, parental status, or protected genetic information. (Not all prohibited bases apply to all programs.) Persons with disabilities who

require alternative means for communication of program information

(Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at 202-720-2600 (voice and TDD).

To file a complaint of discrimination, write USDA, Director, Office of Civil Rights, Room 326-W, Whitten Building, 14th and Independence Avenue, SW, Washington, DC 20250-9410 or call 202-720-5964 (voice and

TDD). USDA is an equal opportunity provider and employer.

Exhibit C

U.S. DEPARTMENT OF AGRICULTURE AGRICULTURAL MARKETING SERVICE SCIENCE AND TECHNOLOGY PLANT VARIETY PROTECTION OFFICE **BELTSVILLE, MD 20705**

OBJECTIVE DESCRIPTION OF VARIETY Wheat (Triticum spp.)

NAME OF APPLICANT (S)

your application.

TEMPORARY OR EXPERIMENTAL DESIGNATION

VARIETY NAME

TX98V9628

ADDRESS (Street and No. or RD No., City, State, Zip Code and Country)

FOR OFFICIAL USE ONLY PVPO NUMBER

PLEASE READ ALL INSTRU	CTIONS CAREFULLY:
Place the appropriate number (0.9)	that describes the varietal character of this variety in the boxes below. Place a zero in the first box (e.g., 0 9 9 or
Comparative data	ss or 9 or less respectively. Data for quantitative plant characters should be based on a minimum of 100 plants
determine plant colors:	ieties entered in the same trial. Royal Horticultural Society or any recognized color standard may be used to
designate system used:delay progress of	Please answer all questions for your variety; lack of response may

1. KIND:										_
1	1 = Common 2 = Durum 3 = Club 4 = Other (Specify)			20	0		0	2	7	Ą.
2. VERNALI	ZATION:			*******				-	•	
2	1 = Spring 2 = Winter 3 = Other (Specify)									
3. COLEOPT	TILE ANTHOCYANIN:									
2	1 = Absent 2 = Pres	ent								
4. JUVENILE	PLANT GROWTH:						-			
2	1 = Prostrate 2 = Semi	i-Ered	ct 3 = Erect							
5. PLANT CO	DLOR: (boot stage)									
3	1 = Yellow-Green									
<u> </u>	2 = Green 3 = Blue-Green									
6. Flag Lea	F: (boot stage)				*					
		! = Re	ecurved					-		
2	1 = Not Twisted 2	= Tw	. بنمغه ط							
2										
<u> </u>		= Wa	ax Present (more on the back of t	he lea	f)					
7. EAR EMER	GENCE:									
1 1 4	Number of Days (Average)	•							
0 5	Number of Days Earlier Th	nan	* <u>Cutter</u>			_				•
	Same As		*							
0 1	Number of Days Later Tha	n	*_TAM 110							
<u></u> _			*Relative to a Commercial Variet	у Gro	wn i	in th	e Sa	me	Tria	J
ANTHER CO	DLOR:									
2	1 = Yellow 2 = Purple			٠						

Exhibit C (Wheat)

9. PLANT HEIGHT: (from soil to top of head	i, excluding awns)		•
0 6 5 cm (Average)			
0 5 cm Taller Than TAM W-1	01		*
Same As TAM 110	and Jagger	·	*
0 2 cm Shorter Than Cutter			*
10. STEM:			**************************************
A. ANTHOCYANIN	D.INTERNODE		
1 1 = Absent 2 = Present	1 = Hollow	2 = Semi-Solid	l 3 = Solid
	4 Number of No	des (average)	
B. WAXY BLOOM	E. PEDUNCLE		
1 1 = Absent 2 = Present	1 1 = Erect	2 = Recurved	3 = Semi-Erect
	3 8 cm Length (av	erage)	
C. HAIRINESS (last internode of rachis)	F. AURICLE		
1 1 = Absent 2 = Present	1 Anthocyanin	1 = Absent	2 = Present
	1 Hair:	1 = Absent	2 = Present
11. HEAD: (at maturity)			
A. DENSITY 1 = Lax 2 = Mid-dense (Laxidense) 3 = Dense	C. CURVATURE 1 = Erect 2 = Inclined 3 = Recurved		
3. SHAPE 1 = Tapering 2 = Strap 3 = Clavate 4 = Other (Specify)	D. AWNEDNESS 1 = Awnless 2 = Apically Awn 3 = Awnletted 4 = Awned	nletted	
2. GLUMES: (at maturity)			
COLOR 1 = White 2 = Tan 3 = Other (Specify)red	E. BEAK WIDTH 1 = Narrow 2 = Medium 3 = Wide		Managain kalif

B. SHOULDER 1 = Wanting 2 = Oblique 3 = Rounded 4 = Square 5 = Elevated 6 = Apiculate 7 = Other (Specify)	F. GLUME LENGTH 1 = Short (ca. 7mm) 2 = Medium (ca. 8mm) 3 = Long (ca. 9mm)
C. SHOULDER WIDTH 1 = Narrow 2 = Medium 3 = Wide	G. WIDTH 1 = Narrow (ca. 3mm) 2 = Medium (ca. 3.5mm) 3 = Long (ca. 4mm)
D. BEAK 1 = Obtuse 2 = Acute 3 = Acuminate	
13. SEED:	
A. SHAPE 1 = Ovate 2 = Oval 3 = Elliptical 4 = Other (Specify)	E. COLOR 1 = White 2 = Amber 3 = Red
B. CHEEK 1 = Rounded 2 = Angular 3 = Other (Specify)	F. TEXTURE 1 = Hard 2 = Soft
C. BRUSH 1 = Short 2 = Medium 3 = Long 1 = Not Collared 2 = Collared	G. PHENOL REACTION 1 = Ivory 4 = Dark- Brown 2 = Fawn 5 = Black 3 = Light- Brown
D. CREASE 1 = Width 60% or less of Kernel 2 = Width 80% or less of Kernel 3 = Width Nearly as Wide as Kernel	H. SEED WEIGHT 30 g/1000 Seed (average) (whole number only)
1 = Depth 20% or less of Kernel 2 = Depth 35% or less of Kernel 3 = Depth 50% or less of Kernel	1. GERM SIZE 2 1 = Small 2 = Mid-Size 3 = Large
4. DISEASE: (0 = Not Tested 1 = Susceptible 2 = Re	esistant 3 = Intermediate 4 = Tolerant)
PLEASE INDICATE THE SPECIFIC	,
2 Stern Rust (Puccinia graminis f. sp. tritici)	1 Leaf Rust (Puccinia recondita f. sp. tritici)
Stripe Rust (Puccinia striiformis)	0 Loose Smut (Ustilago tritici)
Tan Spot (Pyrenophora tritici-repentis)	Flag Smut (Urocystis agropyri)

Exhibit C (Wheat)

Contario avenno (Cranidad I ant Diassa)	warf Bunt (Tilletia controversa)
0 Septoria avenae (Speckled Leaf Disease) 0 Ka	arnal Bunt (<i>Tilletia indica</i>)
0 Septoria tritici (Speckled Leaf Blotch) 2 Po	wdery Mildew (Ersiphe graminis f. sp. tritici)
0 Scab (Fusarium spp.) 0 "S	now Molds"
	ommon Root Rot (<i>Fusarium, Cochliobolus</i> d <i>Bipolaris</i> spp.)
3 Barley Yellow Dwarf Virus (BYDV) 0 RI	nizoctonia Root Rot (Rhizoctonia solani)
	ack Chaff (Xanthomonas campestris pv. Inslucens)
	cterial Leaf Blight (<i>Pseudomonas syringae</i> . <i>syringae</i>)
3 Wheat Streak Mosaic Virus (WSMV)	her (Specify)
Other (Specify) Ot	her (Specify)
	her (Specify)
Other (Specify) Ot	nor (opcony)
	ther (Specify)
Other (Specify)O 15. INSECT: (0 = Not Tested 1 = Susceptible 2 = Resista	ther (Specify) nt 3 = Intermediate 4 = Tolerant)
Other (Specify)	ther (Specify) nt 3 = Intermediate 4 = Tolerant) eeded)
Other (Specify) Other	ther (Specify) nt
Other (Specify) Other	ther (Specify) nt
Other (Specify) Other	ther (Specify) nt
Other (Specify) Other	ther (Specify) nt
Other (Specify) Other	ther (Specify) nt

16. ADDITIONAL INFORMATION ON ANY ITEM ABOVE, OR GENERAL COMMENTS:

Exhibit D

Additional Description - 'TAM 112' Wheat

TX98V9628 is adapted to the rain-fed production systems in the High Plains of Texas and similar areas in adjacent states. It has been extensively tested throughout the Great Plains. It is resistant to the three most prevalent greenbug biotypes (E, I, and K). Based on pedigree and reaction to greenbug, it has the genes gb2 and gb3. Based on field observations, it is resistant to powdery mildew and susceptible to stripe rust. During selection and testing, it was resistant to the prevalent races of leaf rust, but is susceptible to a race that is increasing in prevalence in Texas. Test weight is medium-high and milling and baking characteristics are good. It does have the 1AL.1RS translocation where the short arm of chromosome 1 in rye has been substituted for the short arm of chromosome 1 A in wheat.

TAM 112 is similar to TAM 110 in area of adaptation, drought tolerance, and disease and insect resistance; but averaged over 46 site-years in Texas, TX98V9628 yielded 5.9 bu/a more than TAM 110 (Table 1). Additional information and data is provided below in within the attached Tables.

Grain Yield

Performance data of TX98V9628 is presented for 103 different site-years:

Loc	ations
2002 North Texas Elite (NTE), Emphasis on Blacklands	9
2002 West Texas Elite (WTE), Emphasis on High and Rolling Plains	11
2002 Southern Regional Performance Nursery (SRPN), 10 states	26
2003 Texas Elite (TXE), statewide	10
2003 SRPN	31
2004 Uniform Variety Trial (UVT), statewide	<u>16</u>
Site-Years	103

Yield of TX98V9628 across the 46 site-years in Texas averaged 53 bu/a (Table 1). This was well above the check cultivars TAM 110 (47), Jagger (47) and Ogallala (49) and slightly less than 'Cutter' (54). TX98V9628 appears to have a wide area of adaptation. The SRPN represents 10 states throughout the Great Plains. TX98V9628 ranked 1st/46 across 26 locations in 2002 and 10th/46 across 31 locations in 2003 (Table

Grain yield at individual locations in Texas is shown in Tables 13-16. TX98V9628 has performed well across all regions of Texas, but seems to be particularly well suited to dryland production on the High Plains and Rolling Plains. The 2002 and 2003 yield trials at Claude were grazed until late February and the 2003 trial at Chillicothe was clipped in mid February to simulate grazing. The grain yield for these three site-years indicates that TX98V9628 should perform well in a dual-purpose system. In 2004, TX98V9628 was tested against current popular varieties in the statewide variety trial. It ranked 3rd statewide, 1st in High Plains Dryland, 2nd in the Rolling Plains, 14th in High Plains Irrigated, and 13th in East/Central locations (Table 13).

Test Weight

Test weight from the 2002 to 2003 Texas trials is summarized in Table 2. Averaged across 27 site-years in Texas, TX98V9628 was 3.2 lb/bu greater than TAM 110, 2.2 lb/bu greater than Jagger, similar to Cutter, and 0.4 lb/bu less than TAM 400 and Ogallala. The test weight of TX98V9628 has been fairly stable across locations, consistently ranking among the top performers except at locations where leaf rust or stripe rust was severe. TX98V9628 is susceptible to some prevalent races of leaf rust and stripe rust. In the 2004 UVT, the test weight of TX98V9628 was equal to or better than all other varieties at the High Plains and Rolling Plains locations but was less than Ogallala and Cutter across the four East and Central locations (Table 13). Across 49 site-years in the SRPN, TX98V9628 was 2 lb/bu more than TAM 107 and not less than any of the checks (Table 8). Overall, it seems that TX98V9628 has a very good test weight except when foliar diseases to which it is susceptible are severe.

Forage Yield

Forage trials were conducted at Overton, Texas in 2003 and 2004 (Table 6). TX98V9628 yielded similarly to the forage wheat cultivar Lockett in both years. The fall and winter forage production (cuttings through February) was particularly good in both years, yielding in the top performing group for most cutting dates. This is a good characteristic for a dual purpose (grazing plus grain) wheat variety.

Agronomic data

In Texas, the heading date of TX98V9628 was 1 day later than TAM 110 and Jagger and 5 days earlier than Cutter (Table 3). Plant height was similar to TAM 110 and Jagger (Table 3). Data from the SRPN confirm that TX98V9628 is medium-early maturity and medium height (Table 8). Winter survival notes taken at locations in South Dakota, Iowa, and Nebraska, indicate that the winter hardiness of TX98V9628 is similar to or slightly less than TAM 107 (Table 8). This is adequate for the southern Great Plains. The straw strength of TX98V9628 has been less than TAM 107 and TAM 110 at high yielding environments (see lodging note in Table 8 and 13). This is a limitation for high input irrigated production. Data is limited, but TX98V9628 appears to be tolerant to low pH soils (Table 8).

Disease and Insect Resistance

Seedling inoculations indicate that TX98V9628 is resistant to the three most prevalent greenbug biotypes (E, I, and K; data not presented). Based on pedigree and reaction to greenbug, it has the genes gb2 and gb3 (the same combination of genes is in TAM 110). Natural infestations were severe enough for field ratings at Etter in 2002 (Table 3) and at Bushland dryland in 2004 (data not presented). At both locations, TX98V9628, TAM 110, and other lines/varieties thought to carry the gene gb3 were rated as significantly less damaged.

TX98V9628 is thought to carry the leaf rust resistance gene *lr41*. During selection and preliminary yield trials it was resistant to prevalent races of leaf rust. But starting in 2003 and increasing in prevalence in 2004, leaf rust races virulent to *lr41* have become dominant in south, central, north central, and northeast Texas (ratings shown in Tables 3, 10, and 13). TX98V9628 is resistant to stem rust and is postulated to carry the gene *sr24* (Table 9). TX98V9628 is susceptible to stripe rust (Tables 3 and 10). It is highly resistant to powdery mildew (Table 11). TX98V9628 does have the 1AL.1RS translocation where the short arm of chromosome 1 in rye has been substituted for the short arm of chromosome 1A in wheat. This translocation has been associated with powdery mildew resistance, resistance to biotype C greenbug, and a moderate level of resistance to wheat streak mosaic virus. Data collected from the SRPN confirms the moderate resistance to wheat streak mosaic virus (Table 11). Based on SRPN data, TX98V9628 appears to be

susceptible to wheat soilborne mosaic virus but might have some resistance to barley yellow dwarf virus (Table 11).

Quality

Protein content, kernel size, and kernel hardness are average for hard red winter wheat (Table 4). Averaged over 11 site-years, the protein content of TX98V9628 was 0.4% higher than TAM 110 and 0.8% lower than Jagger. The kernel weight was 0.7 mg more than TAM 110, 2.2 mg more than Jagger, and 2.5 mg less than TAM W-101. The average single kernel hardness (SKHT) value was 72, compared with 70 for TAM 110 and 72 for Jagger. This is acceptable for a hard wheat variety. TX98V9628 graded hard at all 4 locations evaluated in 2003. Mixograph and baking data from analyses performed by the USDA-ARS Hard Winter Wheat Quality Laboratory in Manhattan, Kansas are shown in Tables 5 and 12. TX98V9628 was acceptable for all characteristics measured. It was similar to TAM 110 and TAM 107 for all characteristics except mixograph time, mixograph tolerance, and baking mix-time. The dough mixing time measured by both the mixograph and baking tests were around 2 minutes longer than either TAM 110 (Table 5) or TAM 107 (Table 12). Dough mixing tolerance was better than either TAM 110 (Table 5) or TAM 107 (Table 12). The crumb grain, crumb color and loaf volume all indicated that TX98V9628 was among the best tested for bread-baking quality. Mixograms from three locations in Texas are shown in Figure 1. They graphically illustrate the long dough mix-time combined with good stability of TX98V9628. In summary, TX98V9628 seems to be a strong dough mixing wheat with an average protein content and good baking quality.

Summary

TX98V9628 has a good yield record across a wide range of environments. It is particularly suited for dryland production systems in the High Plains of Texas and similar areas in adjacent states. It should perform in dual purpose (grazing plus grain) systems as well as grain only systems. TAM 112 is similar to TAM 110 in maturity, area of adaptation, drought tolerance, and greenbug resistance. It is better than TAM 110 for grain yield, forage yield, test weight, and milling and baking quality. Like TAM 110, it does have the 1AL.1RS translocation where the short arm of chromosome 1 in rye has been substituted for the short arm of chromosome 1A in wheat.

Table 1. Grain yield data from TAES trials conducted during 2002 to 2004

		Α	Average	# Locations	52.9 46		47.0 46						76
•		2002	NTE Rank/40		53.0 15			45.0 36		50.7 21	1	44.9 37	O
	Grain Yield		WTE Rank/40	(bu a ⁻¹)	47.5 3	37.3 36	42.2 19	41.0 29	43.2 12	43.6 8	42.8 15	41.5 25	7.7
		2003	TXE Rank/40		49.4 16	42.0 38	42.7 36	45.7 31	56.4 1	48.6 22	51.1 10		10
		2004	UVT Rank/40		58.7.3	47.8 24	54.4 15	51.8 19	56.2 11	52.4 18	58.4 6	ı	16
	14	Name			TX98V9628	TAM W-101	TAM 110	Jagger	Cutter	Ogallala	TAM 111	TAM 400	# Locations

UVT = Uniform Variety Trial, Statewide

TXE = Texas Elite, Statewide

WTE = West Texas Elite, Emphasis on High Plains and Rolling Plains

NTE = North Texas Elite, Emphasis on Blacklands

UA1 = Uniform Advance 1

Table 2. Test weight data from TAES trials conducted during 2002 to 2003

) (est weight	tht					
				2003						20	2002	
[TXE						WTE	NTE	Average
Bush Irriga	Bushland Irrigated	Chillicothe	Munday	McGregor Brady		College station		Uvalde	uling Uvalde Average			across years
						lb bu-1						
X98V9628 62.1	1.1	62.0	0.09	61.4	60.4	62.0	57.2	58.9	60.5	59.8	58.8	59.7
	ø.	61.6	59.2	61.1	57.1	61.0	56.2	53.7	58.9	58.5	57,4	58.3
	4.	61.0	27.7	58.7	57.5	57.4	52.5	57.1	57.8	56.7	55.1	56.5
TAM 400 62	62.6	62.5	6.09	63.1	60.7	63.3	57.4	58.5	61.1	60.5	58.7	60.1
	6.	61.8	58.1	59.6	58.0	59.3	56.3	57.4	58.8	57.6	56.3	57.5
	œί	62.3	0.09	61.2	59.4	62.9	59.6	55.8	60.4	59.3	59.1	59.6
62	7	62.4	8.09	61.4	58.8	62.6	58.0	55.2	60.2	60.3	0.09	60.1
62	7	61.4	58.5	60.1	58.6	60.4	56.0	51.8	58.6	59.1	ı	
ocations oc									8	10	6	27

TXE = Texas Elite, Statewide
WTE = West Texas Elite, Emphasis on High Plains and Rolling Plains
NTE = North Texas Elite, Emphasis on Blacklands

Table 3. Agronomic and disease data from TAES trials conducted during 2002 to 2003

ا (د		Ĭ	Heading			He	Height			Dispase	2	Croophy
Name	2003		2002	Average	2003	2002	2	Avorage		2003		2002
	X	WTE	WTE NTE		7XE	WTE NTE	NTE	añe lañe	Leaf rust	Leafrust	Stripe rust	Effer
			doy 1			Ö	cm		Lulina	Castroville	Castroville	(0-5 5=Dead)
TX98V9628	124	118	107	114	72	56	99	65	, S09	MR	S	0.57
TAM W-101	124	124	112	119	65	54	62	09	808	MR	N N	. c
TAM 110	123	116	106	113	71	58	63	64	80S	S	, U,	7 33
TAM 400	124	119	108	115	99	54	64	61	DEAD	WS WS	S N	2.5
Jagger	123	117	106	113	9/	58	63	99	DEAD	v.	MR.	
Cutter	125	126	110	119	74	29	99	67	205	œ	Ĺ	70.5 6
Ogallala	125	110	112	114	99	28	65	63	20MR	: c	AD.	0.00
TAM 111	124	121	ı		74	59) i	<u> </u>	S09	-م :	í a	3.07 3.67
# Locations	2	3	4	6	2	2	_	22				50.0
mean												3 33
cv (%)		;										16.20
l.s.d. (0.05)												0.05

¹ Days of the year from January 1

TXE = Texas Elite, Statewide

WTE = West Texas Elite, Emphasis on High Plains and Rolling Plains NTE = North Texas Elite, Emphasis on Blacklands

²Rust sores: the numeric score indicates % of flag leaf area infected and the infection type is indicated by R= resistant, MR= moderately resistant, MS= moderately susceptible, S= susceptible, t= trace, ?= not scored, Dead= no leaf area to score.

Table 4. Protein and single kernal hardness data from TAES trials conducted during 2002 to 2003. TAMU Wheat Quality Laboratory.

					Quality	litv		
Momo		2-years	2-years Average				2003	
	Protein	SKHT Weight	SKHT SKHT Diameter Index	SKHT Index	Single	Kernel Hardness	Single Kernel Hardness Test (SKHT) Distribution	ion
	mp %	тв	тш		Bushland Irrigated	Chillicothe	Prosper	McGregor
TX98V9628	14.8	29.3	2.0	72	0410285801 0	0004148201	0410285801 0004148201 10713555507 1112138411	102138401
TAM W-101		31.8	2.1	9	0624373302)410206601	0624373302 0410206601 0412255901 102162760-01	716226001
TAM 110		28.6	1.9	20	0821254602 0	1103128401	0821254602 0103128401 0513295301 0003079001	7030701
TAM 400	15.2	27.4	2.0	74	0209216801 0	1101089001	0209216801 0101089001 0512176601 0103069001	103069001
Jagger		27.1	1.9	72	0210226601 0	003138401	0210226601 0003138401 0816225402 0106138001	106138001
Cutter		30.1	2.0	71	0310256201 0	003138401	0310256201 0003138401 0310256201 0101079101	101079101
Ogallala		25.2	1.9	99	0512245901	513245801	0512245901 0513245801 1213264903 0614186202	314186202
TAM 111	14.5	28.2	1.9	61	0519314501 0	721304202	0519314501 0721304202 19-23273103 0920284302	9-20284302
# Locations	11	6	6	6				

			,						į	Σŀ	MIXOGRAPH	핅							B/	BAKING			
Τ̈́e	Test Entry	Мате	Test weight	weight	diameter	index	x distribution	Wheat Ash	Wheat Protein	Flour Yield	Flour F	Flour Mis Protein tir	Mixing Mi time tole	Mixing tolerance ab	Flour	mixing time	Proof height	Loaf C	Crumb (Crumb color	Loaf	Poly phenol oxidase	Brightness after 24 hrs
i			(lb bu ⁻¹)	(mg)	(mm)	Ш		(%)	(%)	(%)	(%)	m) (%)	(min)	(0-61)	(%)	(min)	(ac)	1	(19-0)		(00)	(long or long)	
¥	CM (1	TAM 110	9. 9.	8 7:1	7 K	92	001-005-016-078-01	1.29	12.3	0700	300000		12	2	61.7	2,99	(3)		700	Creamy	(22)	(AU/min/mL	SUPE
11 X	9 <	TAM 400	979	4.00 4.00	2.53	88	000-005-023-072-01	6	e e e	90°3	0.32	10°51	4 2	VI.	ζ <u>ο</u> (Ω	3.08	7.5	149.6	. 44 . 103	Creamy	860	6290	2 2
	T 147	Jacobir Jacobir	64.4 81.6	32.4 39.5	2.41	2 4	001-001-005-093-01	1.42	13.7	- 33	21 101 101	200	3.45	4	64.0	5.23	7.9		4.0	creamy	990	0.260	72.15
TXE	9	Cutter	62.8	34.5	2.47	7 7	000-000-010-01-01	1.00 2.00	15.4	00 P			33	cu c	62.6	3,75	7.4	150,7	ය ප්	creamy	890	0.432	69.43
TXE		Ogallala	63.6	28.1	2 2	75	000-007-010-000-01	 	0.0 0.0 10.0				S, E	თ,	62.7	4.75	9.7	150.5	3.8 8.0	creamy	945	0.543	68.74
X H	. ю	TX96D1073	62.4	32.7	2.38	93	001-005-023-067-01	147	0.47		20.00	7.7	2.50	- c	9.09	9.00	7.7	149.1	4.0	creamy	960	0.530	71.39
꿆	-	TX00D1390	63.2	33.6	2.48	67	001-005-023-071-01	1.38	5 5		•		451	ם ער	98.6 62.3	7.5C 86.0C	7 7	148.4	89 E	creamy	855	0.620	70.78
2	때 r 은 7	TX01D3232	80.8	31.7	2.42	77	002-002-008-088-01	1.36	13.0		•		3.21	ာက	62.6	4.59		150.1	0. C	creamy	855	0.529	72.50
<u> </u>		1X00V111/	22.5	32.6	2.54	72	000-003-013-084-01	1.33	12.5	-	`		3.00	2	62.0	4.61	7.5	149.6	80.00	creamy	880	0.470	69.40
Υ <u>Σ</u>	ПП Л	TX0145936	ο 2.4 4. α	37.5 2.03	2.38	3 8	000-000-002-098-01	1.51	13.7		_	1,5	3.66	4	64.3	4.24		150.0	3.8	creamy	940	0.557	68.77
1 X	_	TX97D5088	62.0	2 6	2.00	8 8	000-004-018-078-01	97.7	3.6	7.80			2.66	. .	61.6	3.03		149.0	4.5	creamy	880	0.574	71.18
光	•	TX98D1158	62.4	34.7	2.55	2 7	002-008-026-064-01	33	4.01		33.0	20 C	3.08	5 4	62.6	00 F		149.4	3.8	creamy	895	0.141	73.22
꼰	E 16 .	TX99A0154	64.0	34.0	2.51	7	001-003-011-085-01	1.42	13.3				3.40	ാന	63.4	5.87 4.61	7.0	150.9	o c	creamy	9 12 12	0.524	71.37
X X		TX00A0181	64.4	38.6	2.62	75	000-001-006-093-01	1.38	13.0		_		3.14	4	62.9	4.04		150.9	ء م م	Creatily	820	0.520	72.63
<u> </u>	Эд Э	1X01V6016	63.6	35.7	2.60	75	000-002-008-090-01	1.42	12.8	-	•		2.14	۴	60.2	3.14		148.4	3.2	yellow	870	0.487	71.59
Y Y		TX00A0174	07.0	20.2	2.40 38.0	8 8	000-001-004-095-01	1.36		_			3.42	4	63.5	4.24	7.2	152.0		creamy	875	0.461	70.39
! !! !!	23 1 Lii	TX00A0391	63.6	36.0	250	75	000-001-007-092-01	رن بر بر	0 0	0 1.07	990	0.9	2.94	4 .	63.4	3.91		149.4		slight yellow	925	0.598	70.64
ΤÆ		TX01A5937	63.6	34.3	2.54	84	000-002-004-094-01	1.35	13.8	_			2.75	4 <i>c</i>	62.6 62.4	9. 8 9. 8	7 6	151.0	4.0	creamy	840	0.492	70.63
뽔!	77	TX01V5314	60.4	34.1	2,50	11	000-004-007-089-01	1.36	14.2	_	_		3.88	ΙN	63.5	4.50		151.7	4.0	Creamy	870	0.602	10.41
¥ 2	, 0 6	TX01V5/19	63.6	35.2	2.65	29	002-005-020-073-01	1.38	14.0		-	9	2.88	_	64.3	3.75		150.9	5.0	creamy	935	0.153	72.94
불	. 72	TX01112527	62.2 8.08	 	2.50 5.50	9 9	000-003-007-090-01	1.37	13.7		_ ,		3.00	~ .	63.9	4.50		150.3	5.0	creamy	950	0.467	70.34
뿚	8	TX0102598	62.4	32.5	2.47	7 4	001-003-008-088-01	S S		69.C	0.30	17.8	3.57	4 4	65.3	5.16		148.1	8.4	creamy	1000	0.489	70.06
뿚		TX01U2723	63.6	31.1	2.40	23	000-000-004-096-01	1.36	12.6	_	-		2.66	1 4	64.4	5 4	7.0 7.0	15.0	4, r. U, C	creamy	040 000	0.587	71.63
뿚		TX01U2733	63.6	33.0	2.45	80	000-001-004-095-01	1.45	13.1	_			2.71	. 4	63.4	4.29	•	150.5	5.0	Creamy	875	0.658	71.75
Y \		1X0102685	64.4	28.5	2.23	72	000-002-012-086-01	4.	14.5	_			3.50	~	63.2	4.00	•	148.6	4.2	creamy	985	0.550	70.56
	3 8	TX95\/4339	63.0	33.3	5, 5 5, 5 5, 5	3 25	002-004-019-075-01	1.37	14.3 6.4		•		3.50	4	64.6	4.25	7.7	150.3	4.0	creamy	940	0.483	71.83
X X	.45	TX00D2234	62.0	33.5	2 42	- 72	001-004-016-079-01	74.	4 c 7 t	09.60	1.50		3,13	24.0	63.7	4.75		149.2	3.5	creamy	925	0.445	71.82
Ž.	8	TX01D3215	64.8	35.5	2.70	2	000-003-015-082-01	39	. 4		- ,-	2.4	250	v -	62.0 62.9	2.55 2.75	. , . ,	149.1	د د ∞ د	yellow	860	0.458	70.50
본	Ж	TX01D3218	63.6	33.4	2.45	77	000-001-009-090-01	1.35	12.7				2.17	_	62.8	2.86	٠	151.3	9 00	creamy	850	0.048	72.27
	37	TX01D3472	63.6	33.6	2.47	33	001-002-010-087-01	.38	13.8	_	Ξ.	· •	5,47	. 9	63.7	71.7	•	150.7	3.5	creamy	006	0.586	70.93
2 2	გ წ	1X99A0556	25.5	30.1	2.53	2 5	000-001-012-08/-01		12.6		•		3.60	4	63.4	5.14	7.5	150.7		yellow	860	0.555	71.55
본	3 4	TX01V6334	62.8	40.5	2.76	r 69	000-002-016-087-01	13,42	4.6	0.00	0.33	0.17 0.47	5.70 2.07	4	67.5	5.24	7.7	151.4	3.0	slight yellow	815	0.632	69.27
55	2.2	TX98D1170	0.5	34.6	2.50	98	003-003-018-076-01	134	12.4	679	48.			· CI	90.8	2 % ; i & :			2,2	creamy	783	0.469	70.27
3	29	TX01M5009	90.8 60.8	27.7	2.24	82	000-007-013-085-04	1.43	13.7	95.7 U	8 8 L L	1.9 4. 1.6 1.	4.32 1.42	4 ~	63.6 60.6	4.93 2.02	7.6 6.8	150.8 152.5	3.5 1.0	creamy	945 685	0.589	69.42 70.04

SKHT= Single Kernel Hardness Test

*0= worst, 6= best

						Forage Yield (Dry matter)	'ield (Dr.	' matter)						
			2002-03	1		0			2003	2003-042				
Nov 7	Feb 11	Mar 10 Mar 26	Mar 26	Apr 29	Total	Rank/34	Dec 4	Jan 29	Mar 10	Apr 16	Total	Rank/30	2-years average	verage
							·lba~¹							Rank
1438	1601	932	714	225	4909	6	730	1530	1604	3079	6943		5926	
1551	1710	268	636	428	5221	7	746	1469	1917	2177	6309	. 2	5765	٠ ،
2007	880	1137	1023	325	5329	5	815	902	1983	2455	6155	5	5742	ı က
1236	1432	1015	534	593	4800	13	744	1502	1678	2227	6151	9	5476) 4
1069	270	1375	950	1225	4889		899	633	2182	2560	6043	œ	5466	LC.
1521	1176	1011	615	625	4947	GO.	849	1461	2220	1290	5820	, 7	25.5) (C
1484	1512	855	524	463	4837	12	556	1363	1569	2030	5518	2	5178	<u> </u>
1331	1132	1011	661	472	4606	18	516	1066	1875	2284	5741	5	5174	- 00
959	1206	980	540	545	4229	29	473	1713	1936	1833	5955	· 	5092) O
1507	1884	645	408	410	4854	=	761	1423	1956	1155	5295	23	5075	3 0
1500	1110	904	410	460	4383	23	548	1036	1697	2454	5735	16	5059	-
1267	635	096	746	468	4076	31	424	1199	1807	2445	5875	13	4976	12
1375	1124	1013	532	348	4392	22	533	1302	2159	1190	5184	25	4788	<u>.</u>
1349	902	982	299	176	4075	32	512	1166	1405	2257	5340	21	4708	4
1049	632	966	724	368	3768	34	883	1148	1325	2192	5548	19	4658	5.
1403	906	1091	536	284	4219	30	262	1006	2035	1253	4891	29	4555	16
1328	1243	921	672	544	4705		656	1305	1878	1864	5703			
26	33	55	53	47	16		238	228	257	271	494			
311	375	188	144	231	693		40	19	15	16	ග			

¹Planted on September 13, 2002. Fertilizer: Preplant 91 lb N, P₂O₅ and K₂O/a. Topdressed with 40 lb N on November 12, 33 lb N/a on January 21, 40 lb/a of N, P2O5 and K2O on March 6 and 40 lb N/a on April 15, 2003. ²Planted on September 15, 2003. Fertilizer: Preplant 700 lb 13-13-13/a. Topdressed with 40 lb N/a on December 10, 2003, 25 lb N/a on February 19, 2004, and 42 lb N/a on March 31, 2004.

³TX95-38-2, TX95-24-1, TX95-16-1, TX95-118, TX97-172, TX89-81-1, and TX95-115 are soft red winter wheat experimental lines from Lloyd Nelson's program. The others are all hard red winter wheat lines or varieties.

		20	2003			20	2002		2-year	2-years average
Name	State		Region				Region			
	Average	Average Rank/46 Average Rank/46	Average	Rank/46	Average	Average Rank/46	Average	Average Rank/46 State	State	Region
						Yield (bu a ^{.1})	a'1)			
TX98V9628	62.2	21	68.8	10	54.5	2	51.3	_	58.3	60.1
Kharkof	40.4	46	43.5	46	28.5	46	27.1	46	34.5	35.3
Scout 66	54.7	43	52.1	45	39.5	45	35.0	45	47.1	43.5
TAM 107	59.1	36	6.09	42	46.7	33	42.3	30	52.9	51.6
Trego	64.9	7	67.2	18	51.1	10	45.3	7	58.0	56.2
# Locations	3		31		4	17.7	26		7	57
mean	61.0		65.4		48.2		42.6			5
cv (%)	11.0		11.5		10.0		13.0			
l.s.d. (0,05)	10.5		43		0		7 %			

Table 8. Mean agronomic data from Southern Regional Performance Nursery (SRPN) conducted from 2002 to 2003

pic	- Ge	1RS		Je)	1AL.1RS	NON.1RS	NON.1RS	1AL.1RS	NON.1RS	2
Average Acid	soil tolerance	(1-5;	1=tolerant,	5=susceptible)	2.4	4.1	3.9	4.0	4.0	2
Average	survival	% stand			63	85	83	70	80	4
Average	Lodging	%			20	70	70	25	20	2
Average	Height	cm			80	100	9	9/	75	32
Average Average Average	Heading	¹ doy			133	143	138	132	137	27
Average	ΛΛ	lb bu ⁻¹				58.7	59.4	27.8	60.1	49
	11	Name			TX98V9628	Kharkof	Scout 66	TAM 107	Trego	# Locations

¹ Days of the year from January 1

Table 9. Stem rust data from Southern Regional Performance Nursery (SRPN) conducted from 2002 to 2003

													٠.
22						·	Stem Rust						
S A			2003						20	2002			
() V	Stem Rust	Seedling	Seedling	Seedling	Seedling	Stem Rust	Seedling	Seedling	Seedling	Seedling	Seedling	Seedling	Seedling
Ivalue		reaction to	reaction to	reaction to	reaction to		reaction to	reaction to	reaction to	reaction to		reaction to	reaction to
	(St Paul,	Stem Rust	Stem Rust	Stem Rust	Stem Rust	(St Paul,	Stem Rust	Stem Rust	Stem Rust	Stem Rust		Stem Rust	Stem Rust
		isolate (St	isolate (St	isolate (St	isolate (St		isolate (St	isolate (St	isolate (St	isolate (St		isolate (St	isolate (St
	QIN)	Paul, MN)	Paul, MN)	Paul, MN)	Paul, MN)	MN)	Paul, MN)	Paul, MN)	Paul, MN)	Paul, MN)		Paul. MN)	Paul MN)
	adult/field	01-TX 27C	74-MN	72-MEX	668 NM-69	adult	NM-08	72-MEX	98-EGY 5B		_	01-TX 27C	
			1409A										
		1111	TPMK	53A RTQQ	QTHJ		633B TPMK	53A RTQQ	RRTS	QTHJ	PTHS	1111	gene(s)
TX98V9628	20 MR	+	+	↽	+	5R	2=	2=	1	2=	2=	2=	24
Kharkof	30 MRMS	4	4	34	23+	S09	တ	တ	03	ဟ	S	S	· ~
Scout 66	10 MRMS	33+	3+	Z + -	2+3+	TMR	S	• -	ഗ	7	ဟ	တ	17
TAM 107	30 MR	7	_	1/3	22-	809	0	-	5=	2=		<u>2=</u>	6.17.24
Trego	20 MR	22+	1-	~	22-	30MR	0	2=	ŗ.	-		2=	6,24

2=small uredinia surrounded by chlorosis; 3=moderate size uredinia without necrosis or chlorosis; 4=large uredinia without necrosis or chlorosis; +=uredinia larger than normal; -=uredinia smaller than normal. A range of infection types is indicated by more than one infection type, with the predominant Seedling infection types: 0=immune response, no sign of infection; ;=hypersensiteve chlorotic or necrotic flecks; 1=small uredinia surrounded by necrosis; type listed first.

Field Rust sores: the numeric score indicates % of flag leaf area infected and the infection type is indicated by R= resistant, MR= moderately resistant, MS= moderately susceptible, S= susceptible, t= trace, ?= not scored, Dead= no leaf area to score.

Table 10. Leaf and Stripe rust data from Southern Regional Performance Nursery (SRPN) conducted from 2002 to 2003

Stripe	2003	Stripe Rust		(Corvallis,		(a)	adult/field	3	ď	Ω	1 v.) V.	ഗ
		Leaf Rust		(Wichita,		(V)	(1-9)		-	٠ (۲	o co	000	က
		Leaf Rust		(Stillwater,		Š	seedling		œ	· v	တ	Ś	N S S
	2002	Leaf Rust		(Stillwater,		Š	adult (1-9)		1	m	9	တ	· -
		Leaf Rust		(St Paul, (Castroville (Stillwater,		Ŷ.	adult		TMS	808	80MR-MS	100S	0
		Leaf Rust				Î	adult		20R-MR	809		S09	208
ust		Seedling	reaction to	Leaf Rust	isolate (St	Paul, MN)	97 CAN 64-	1 KFBJ	ö	ო	2+/3+	<u>o.</u>	ó
Leaf Rust	·	Seedling	reaction to	Leaf Rust	isolate (St	Paul, MN)	99 GA 86 97 CAN 64-	3 TNRJ	3+	3+	3+	3+	• -
		Seedling	reaction to	Leaf Rust	isolate (St	Paul, MN)	00SD520	MCDS	+	3+	3+	ĊΪ	÷
	2003	Seedling	reaction to	Leaf Rust	isolate (St	Paul, MN)	885QN66	THBJ	• -	33+	33+	33+	
		Leaf Rust		(Stillwater,		OK)	seedling		œ	S	တ	တ	MR
		Leaf Rust		(Brookings,		SD)	adult/field		MS	ഗ	တ	တ	2
		Leaf Rust Leaf Rust		(St Paul,		MN)	adult/field		20 S	30 MS	20 MS	20-30 MS	5 MR
- 17		Name	D. COL	:					TX98V9628	Kharkof	Scout 66	TAM 107	Trego

chlorosis; +=uredinia larger than normal; -=uredinia smaller than normal. A range of infection types is indicated by more than one infection type, with the Seedling infection types: 0=immune response, no sign of infection; ;=hypersensiteve chlorotic or necrotic flecks; 1=small uredinia surrounded by necrosis; 2=small uredinia surrounded by chlorosis; 3=moderate size uredinia without necrosis or chlorosis; 4=large uredinia without necrosis or predominant type listed first.

Field Rust sores: the numeric score indicates % of flag leaf area infected and the infection type is indicated by R= resistant, MR= moderately resistant, MS= moderately susceptible, S= susceptible, t= trace, ?= not scored, Dead= no leaf area to score.

Table 11. Other fungal and viral disease and Hessian fly data from Southern Regional Performance Nursery (SRPN) conducted from 2002 to 2003

į	Hessian fly	2002	Hessian fly	•	(Manhattan,		(3)	2		U) U) (o 0	ηI
	Hessi	2003	Hessian fly Hessian fly	•	(Manhattan, (Manhattan.		ίν.	(2)		U	טכ) () (οI
			Wheat	streak	mosaic	Virus	K K	(1-9)		4	rα) h	- د	o 4
			Wheat	streak	mosaic	virus (Hays,	. G	(1-9)		4	۰ ۲	- 1-	یا ۔	າທ
			Barley yellow	dwarf virus		(Columbia,	(OM	%flag leaf	acitorologaip	4	- 52	ر د ر	2 ∘	o €
į		2002	arley yellow		dwarf virus		Urbana. II.)	% Dwarfing		0.0	18.4		2 0	10.9
	sse		Barley yellow Barley yellow Barley yellow		dwarf virus		(Urbana, IL) (Urbana, IL) (Urbana, II.)	(6-0)		4	· (c	ນ	۸ (· ' ' ' ' ' ' ' ' ' ' ' ' ' ' ' ' ' ' '
	Viral disease		Wheat B	soilborne		mosaic virus	Urbana, IL)	(6-0)		6	: 00	α	α	4
			Wheat	soilborne	mosaic virus	(Stillwater, T	Š	(4-1)		2	5	i (r)		ı —
			Wheat	soilborne		mosaic virus	(Wichita, KS	(1-9)	•	8	တ	o.	ο	2 0
		2003	Wheat	soilborne		mosaic virus n	(Winfield, KS (3	ო	က	· cr	1
			Wheat	soilborne	mosaic virus	(Stillwater, P	OK) ((14)		2	_	7	-	1
			Powdery	Mildew		, vvoster,	OH)	(1-9)		_		7		4
	sease	2002	Tanspot		(Winfield,		KS)	(1-9)		4	Ŋ	7	7	9
	Other foliar fungal disease		Septoria				MO)	%canopy	brown		36	20	17	11
	Other foli	က	Septoria	:	(Columbia, (Columbia,		MO)	%canopy	brown	32	33	32	30	26
		2003	Septoria	(West	i afovetto	raicyana,	(<u>,</u>	(1-9)		7	9	7	^	7
	Sr	-{		Name	-					TX98V9628	Kharkof	Scout 66	TAM 107	Trego

Higher numbers (scores and percentages) indicate more severe symptoms

Table 12. Baking and mixograph data from 2002 Southern Regional Performance Nursery (SRPN). USDA-ARS Hard Winter Wheat Quality Laboratory.

ļ		Chemica	<u></u>			Mixograph			Bake	
Name	^	wheat	¥	flour	Water	-	-	Water		- Jag
	Protein Milling	Milling Yield	Ash	Protein	Absorp.	Feak time Iolerance	lolerance	Absorp.	Mix time	Volume
	(%)	(%)	(%)	(%)	(%)	(min)		(%)	(min)	(00)
Composite of South Central P	South Ce	ntral Plains								
Kharkof	14.4	60.2	0.49	12.6	64.0	4.13	4	63.7	4,2	070
Scout 66	13.5	9'29	0.36	12.0	64.0	2.75	- 4		5.5	י ני פי פי
TAM-107	10 x	878	98.0	110	6. 10) i	+ ~) i), r	000
	7.7) †	5.5		7.10	3.88	4	62.0	9.00	950
lrego	13.1	66.4	0.36	1.1	62.5	3.00	7	62.1	4.00	855
TX98D1170	13.6	0.99	0.41	12.0	63.4	3.50	4	62.8	4.75	875
TX98V9628	13.1	65.1	0.39	11.7	63.4	5.50	5	65.3	8.50	960
))
Composite of Southern High P	Southern	High Plains								
-	(]	;							
Knarkot	16.6	27.7	0.49	14.3	66.2	3.75	က	64.2	5.13	895
Scout 66	15.6	2.99	0.38	14.1	65.5	2.38	က	61.8	3.38	930
TAM-107	14.2	64.4	0.36	12.6	63.9	3.00	4	63.0	4.25	905
Trego	13.9	66.1	0.36	12.5	63.8	2.00	7	8.09	3.25	855
TX98D1170	14.7	65.1	0.44	13.5	64.6	2.50	ო	63.9	3.25	096
TX98V9628	14.7	62.1	0.40	13.3	64.5	4.00	4	65.6	00'9	975

2004 Uniform Variety Trial (UVT)- Yield and other ancillary data by region of the state Table 13.

2		Ϋ́	Yield (bu a ⁻¹)	a-1)		-		Test Weight (Ib bu-	siaht (I.	0 bu-1)		E	Ī	Shattering	lodging		-	Conf runt	
)					-				1		T			8	ĺ		רמט	าตอเ	
NAME														0-2	ဝ-ဂ				
9		State	윺	웊		East/			웊		East			0=none	0=none				
	Rank/40	Avg	п	Dry	д Д	Cen	Avg	H III	Irr Dry	윤	Cen	DOY1	£	2=100%	5=100%	PRO	MCG	CAS	GRD
TX98D1170	1	62.5	83.1	32.9	64.5	65.0	56.7	56.9	53.2	61.4	57.6	116	75	0.3	1.8	tMR	c	30R/MR	SONAR+
TX98V9628	က	58.7	79.6	35.1	64.0	52.3	59.2	60.4	56.6	62.4	57.3	113	78	0.4	3.4	1005	1000	3000	1000
TAM W 101	24	47.8	77.9		49.1	31.6	58.0	59.2	56.5	58.4	57.0	110	2 6		† o	300	200	200	1003
TAM 440	7				1) L	1 0) l	5 6		2 :	7	t :	 O	200	202	2001	2001
	2		ο 	4.00	37.5	59.5	5/.p	28.0	55.5	60.7	54.9	113	79	0.0	2.3	100S	100S	100S	100S
Jagger	16		76.9	30.3	54.4	40.0	57.1	58.6	54.8	61.3	56.1	114	92	1.1	2.9	100S	1008	1005	100S
Ogallala	92		77.7		53.2	47.0	58.9	60.2	55.9	61.8	59.2	119	69	1.7	6.0	20MR	50MS	SM09	30MS/MR
Cutter	£	56.2	74.8	31.2	60.1	55.1	59.0	59.0	56.9	62.0	59.5	118	62	1.8	3.4	20S	808	1005	808
TAM 111	9	58.4	90.4	90.4 29.8	56.6	48.5	58.3	58.6	56.1	62.1	58.7	118	78	0.3	5.7	809	808	100S	1005
TAM 202	23	49.1	65.5	29.3	58.4	41.5	57.6	59.0	55.4	61.3	54.8	115	74	3.0	<u>+</u>	808	1005	1005	1005
Sturdy 2K	27	43.8	47.5	25.2	52.2	51.7	57.3	57.1	54.2	6.09	59.2	120	81	3.8	1.	10MR/MS	50MR	40R/MR	60MR/MS
Locations		16	2	4	က	4	14	4	4	2	4	ဖ	-	2	3				

HP Irr= High plains Irrigated (Bushland, Etter, Nazereth, Daihart, and Clovis)

HP Dry= High plains Dryland (Bushland, Etter, Claude, and Clovis)

RP= Rolling plains (Chillicothe, Chillicothe (Hardeman Grain), and Abilene)

East/Cen= East/Central (Prosper, Ellis County, McGregor, and Brady)

DH= Days to Heading (1 Days of the year from January 1)

Ht= Plant height in Centimeter (cm)

Shattering score was taken at Bushland Irrigated and Etter Irrigated.

Lodging score was taken at Bushland Irrigated, Etter Irrigated, and Nazereth. $^{\dagger}A$ few plants were found with susceptible reaction

² Field Rust sores: the numeric score indicates % of flag leaf area infected and the infection type is indicated by R= resistant, MR= moderately resistant, MS= moderately susceptible, S= susceptible, t= trace, ?= not scored, Dead= no leaf area to score.

											,					ċ		2	
Bushland Irrigated	Claude	Chillicothe Munday	Munday	Prosper	McGregor	Brady	College Luling Uvalde	uling		Height Heading	guipe	Leafrus	rust	Stripe rust	Test	ę.	SKHT	SKHT	SKHT
┦╿			Yield	d (bu a')				$\left\ \cdot \right\ $		lis	doy	Luling	Castroville	Castroville		(dm) %		Diamica	
0	25.0		25.1	Ш		52.0	51.0	21.0	1_	-	٦			91		(IIII) e			
94	28.5	513		6	612	200000	0000000				5274		رن ان	Y Y	98.9	6 5 8 6	32.2	2.1	9 8
88.6	22.8	52.4		53.7		58.1	1 1	26.8	1		1		3	2	28.6	14.5	28.4	10	2 K
80.1	16.9	40.4	28.2	41.5	46.3	9.99	49.5	27.2	46.0 65	\dashv	118 DEAD		MS	NS.	61.1	15.1	26.5	6.	73.5
76.6	28.0	42.5	40.4	55.8	64.1	56.5	40.5	28.4	23.7	76	118 DEAD		S	MR	58.8	15.7	27.4	1.9	70.5
89.5	26.5	52.0		43.9	73.1	55.8	61,3	58.0	64.2	73	120 208		œ	·«	60.4	15.3	29.9	2.0	72.3
84.3	20.2	45.3	32.0	54.0	58.3	53.6	51.1	42.4	45.3 65		120 20MR			MR	60.2	15.6	24.4	1.8	63.3
86.8	16.8	49.7	34.0	54.3	52.4	53.0	44.4	18.5	31.2 69	-				MS	56.4	15.1	25.5	- 89	62
	23.0	45.1	32.6			Ġ	59.6		55.0 71		45 .		х	NA.	87.8	2 P	28.2	S	9
2	21.2	38.2	2	54.3	10000	55.0	9	33.3	×		121 20MR		R	αc	59.3	15.0	27.9	1.9	64.3
	952				53.5	60.7	57.3	g4 7	313 72		118 605			v)	60.5	14.5	19 19	2.0	760
	25.3	46.8	27.9		55.0	60.1	56.5	37.4	39.5	\dashv	119 60S		ď	S	6.03	14.3	28.7	2.0	69.8
	27.1	47.6		61.4	64.9	54.0	59.0	45.2	50.5 68		120 5R			œ	60.2	15.3	30.5	2,1	52.5
	6.8	38.5	33.2	50.7	68.1	55.7	58.1	44.4	52.9 69	\dashv	123 5;			œ	59.9	14.5	25.9	- 8.	67.3
77.0	24.5	35.8	35.2	38.3	72.1	57.4	59.8	51.4	9.0	-	117 40;/5R			ıκ	63.5	15.4	31.8	2.1	7.0
86.1	25.3	46.9	30.5	58.2	54.5	52.8	53.1	26.1	40.0 68	-	120 40;			MS	58.1	14.3	27.5	ر ي	61.5
84.0	22.7	46.6	31.8	56.8	64.2	56.1	54.5	35.4	48.2 68	\dashv	119 30;	**	X.	œ	59.2	15.1	27.3	2.0	68.3
105.7	24.9	50.1	27.7	47.7	43.7	46.7	55.6	32.6	40.6 76	4	123 5MR	~		æ	58.5	14.8	25.4	6:	61.5
88.1	23.3	45.4	37.7	58.9	80.6	9.9	71.1	62.1	14.4 66	\dashv	119 tR		~	a.	57.6	14.8	26.7	2.0	64.3
	21.7	43.5	34.6	63.1	74.5	4.4	64.7	61.1	46.0 68	\dashv	120 tR			œ	58.7	14.8	26.0	2.0	69.0
78.5	28.4	46.6	34.2	48.8	68.9	61.9	64.9	26.0	49.7 72		118 5R			MS	6.09	14.2	28.7	2.0	58.0
	22.3	47.9	37.4	51.6	59.3	50.7	59.1	144.1	32.2	\dashv	119 20MR		œ	MS	58.8	14.2	30.1	2.1	59.8
- 1	24.2	44.4	37.5	55.7	63.7	56.6	66.4	39.4	42.3 68	\dashv	119 40M	40MR/MS N	MR	S	61.1	14.8	28.0	1.9	62.3
	22.0	48.3	29.7	58.5	56.9	62.4	62.0	39.2	45.4 79	-	118 30MS	2		VS	61.1	14.3	28.6	2.0	71.5
- 1	20.6	50.2	37.7	58.3	75.8	59.3	69.3	44.2	46.4 69		120 10R	α		œ	59.7	14.5	26.8	1.9	90.5
	28.4	50.9	36.4	54.7	65.4	58.1	64.6	45.9	45.0 69	\dashv	118 5MR	~		MS	58.4	14.5	27.5	2.0	71.8
	25.3	46.8		53.5	55.3	55.9		27.7	49.9 72	-	120 80S		~	MR	59.7	15.0	29.3	2.0	62.8
	26.4	48.3		46.8	53.3	46.3	54.6		40.9 71	\dashv	122 0-20MR	T	MR	MR	58.3	15.4	25.8	8.	65.8
76.7	22.3	44.0		48.6	59.4	54.9	- 1	38.1	48.9 71		119 40MS	ည		MR	59.9	14,8	25.9	1.9	64.3
82.3	30.2	45.7	\perp	45.8	65.4	58.7	64.5	55.1	56.4 68	-	121 tR			S	59.7	14.3	27.8	1.9	52.5
71.0	19.2	43.8		53.8	32.8	46.1	[_		- 1	-	120 0,>STP	A die		S	57.1	15.1	27.3	1.9	76.5
87.7	24.5	46.8		42.3	31.4	46.7	39.8	27.6	27.6 73		122 0,>STP	R R		٧S	60.5	15.6	26.8	2.0	899
70.5	23.6	45.7	ᆜ	51.3	8.69	59.6	64.7	64.8	19.5 79	+	117 fR			MR	59.4	14.1	29.0	2.0	28.8
80.4	24.2	47.0		43.0	59.0	63.5	55.4	49.1	24.5 72	+	117 60S/	60S/10R A	MR	S	61.6	15.1	30.6	2.0	62.8
96.5	22.9	40.9	32.7	46.9	47.4	56.9	48.0	34.2	12.9 68	\dashv	118 DEA	DEAD LF R		vs .	59.2	14.0	25.1	1.8	61.0
92.3	24.5	46.4	29.7	42.7	. 44.7	15.8	56.9	29.6:	43.7 72		118 50M	50MR/MS N	MR	2	58.9	14.4	30.6	2.0	70.3
95.4	23.5	47.1	27.9	50.3	54.3	51.8	56.2 3	30.8	32.5 71		121 50S		MR	MR	60.7	14.5	28.4	1.9	71.3
97.9	22.9	45.6	30.5	41.9	58.6	51.7	53.8	28.6	39.1 71	-	121 10R/S	S) R		MS	58.5	14.1	27.2	9:1	61.3
95.4	25.4	42.1	27.7	47.2	35.3	50.4	42.8 2	24.9	36.7 72		120 10R.	10R,>STP		8	60.5	14.1	30.8	2.0	74.3
88.4	22.9	49.2	37.5	48.0	67.1	1.1	66.5	52.8	52.3 78		119 40M	40MR/MS N	MS	R	61.8	15.0	30.2	2.0	65.5
		;	;	;	1	;					2				6 0	4	4	4	4
85.7	233	45.7		51.1	58.3	ري 6.		.,	39.5										
e Cu	24.8	8.0		හ. ල	5.3	5.0			το το										
4.9	9.1	6.0	38	7.7	2.5	42	67	*											

Table 14. 2003 Texas Elite (TXE) by location

Brady: severe deer damage to awnless (19, 20, 36) Uvalde: freeze damage

Field Rust sores; the numeric score indicates % of flag leaf area infected and the infection type is indicated by R= resistant, MR= moderately resistant, MS= moderately susceptible, S= susceptible, t= trace, ?= not scored.

Dead= no leaf area to score.

Height Uvalde 42.8 South Texas McGregor 46.4 10.5 8.0 Brady 48.8 Prosper 45.6 4.7.5 4.7.5 6.7 54.2 10.9 9.6 Lockett Munday Rolling Plains 48.7 Chillicothe 39.2 15.3 9.7 Etter Dry 9.4 20.0 2.1 Claude 29.7 8.7 4.2 High Plains Bushland Dry 23.7 5.8 2.2 Bushland Irrigated 77.4 6.7 8.5 Average Rank State Sturdy 2K TX95A3091 AP502-CL TX98A5421 TX97A0122 TX97A0122 TX97A0169 # Locations Mean CV% LSD(.05) TAM W-10
TAM 202
TAM 302
TAM 400
Jagger
Ogaliala
2145
Above
Custer 2002 WTE


Table 15. 2002 West Texas Elite (WTE) by location

28

Č				High									•			-			
2002 State Colling plains	plains	plains			plains	I	_	Dallas	- 1	တိ	South Texas	as	Agro	Agronomy		U	Quality		
NTE Average Rank Bushland Chillicothe Munday	Rank Bushland Chillicothe Munday	Rank Bushland Chillicothe Munday	Bushland Chillicothe Munday	Chillicothe Munday	Munday		Prosper	Cooke County	Ellis County	Brady	Brady McGregor	Uvalde	Height	Heading	TW	Protein	SKHT	SKHT	SKHT
				Ш	Ш			Yield (bu a ⁻¹)	J a 1				등	óop	lb bu	%dm	Ē	mm	
01 39.9 40	40 79.7 38.5	79.7 38.5	38.5	38.5	AVANA	į.	31.9	31.2	46.4	42.2	31.2	43.7	62	112	57.4		32.9	2.1	59.5
42.4 39 75.1 41.1	39 75.1 41.1	75.1 41.1	41.1	41.1			64.4 45.1	34.9	48.9	43.1	30.3	48.2	83	106	55.1	7 0 y CO -	30.8	2.0	69 5
45.1 35 82.4 38.7	35 82.4 38.7	82.4 38.7	38.7	38.7	48.7		48.6	41.8	53.5	39.8	32.3	44.9	99	111	55.5	_	29.5	200	714
00 44.9 37 81.7	37 81.7 39.2	81.7 39.2	39.2	39.2	62.7		44.0	35.8	47.5	43.4	27.2	52.1	64	108	58.7	15.3	28.9	2.1	74.2
45.0 36 66.9 40.7	36 66.9 40.7	66.9 40.7	40.7	40.7	59.6	;;, ,	50.8	42.9	48.6	41.5	24.9	41.7	63	106	56.3	-	28.3	2.0	74.5
21 85.5 39.5	21 85.5 39.5	85.5 39.5	39.5	39.5	56.	S	51.9	48.4	55.4	38.2	20.1	59.6	8	112	0.09	15.6	27.5		71.0
31.3 1/ 09.1 44.8 21.0 75 76 40.2	25 76 A0.2	76.6 44.8	44.8	44.8	28	ه اد	55.6	51.9	65.0	41.9	40.3	43.8	200	106	57.6	\rightarrow	28.3	\Box	69.2
58.5 3 81.5 49.6	3 815 496	81.5 49.6	40.6	40.6	5 79	ماد	1 7 7 7 7	- 0.0	92.7	7 9	8.24 0.03	27.7	38	10,7	29.1	13.1	33.4	十	00.1
01 47.0 29 81.4 45.4	29 81.4 45.4	81.4 45.4	45.4	45.4	54.5	1	47.2	44.8	59.3	38.2	31.4	38	8 2	300	57.2	-	ى ن ت ت	- 60	0.0
49.8 24 48.1 33.0	24 48.1 33.0	48.1 33.0	33.0	33.0	54.	m	50.9	54.4		43.9	49.7	40.1	2	11	57.7	-	283	T	83.5
45.2 33 75.2 37.4	33 75.2 37.4	75.2 37.4	37.4	37.4	60.7	-	43.5	36.8		43.6	39.0	48.8	26	107	57.2	_	29.9		79.5
50.8 20 86.2	20 86.2 41.4	86.2 41.4	41.4	41.4	6	=+	57.9	55.4		46.3	38.6	31.4	61	108	61.0		29.7	T	79.2
51.3 18 77.1 34.4	18 77.1 34.4	77.1 34.4	34.4	34.4	62	너	51.9	52.3		48.0	40.8	63.7	99	107	57.3		31.7	1	747
62.3 1 78.5	78.5 44.8	78.5 44.8	44.8	44.8	99	2	99	65.2		51.8	60.1	62.9	65	108	2.09		32.3	2.2	55.8
92.9 44.7	6 92.9 44.7	92.9 44.7	44.7	44.7	61.	ي ا	59.1	56.7		38.0	51.0	38.4	99	110	57.4	H	29.9	П	68.6
5.45 S S S S S S S S S S S S S S S S S S S	8 000 355	37.3	37.3	37.3	o c	N v	186	67.1		50,5	52.5	59.0	0 8	108	56.9	#2500 C	33.6		27.8
53.6 11 68.5 37.2	11 68.5 37.2	68.5 37.2	37.2	37.2	8 6	- 4	540.7	1.00		40.0	53.0	59.5	69	111	58.7		30.6	十	68.4
55.5 5 85.9 43.4	5 85.9 43.4	85.9 43.4	43.4	43.4	63.	- 6	47.9	59.2	62.2	40.5	56.6	56.7	- 6	1 2	2.86.5	+	33.0	2.3	0.40
55.2 7 76.9 43.4	7 76.9 43.4	76.9 43.4	43.4	43.4	57.	7	57.5	58.7		44.9	51.7	54.0	99	108	59.9	15.1	30.4	2.2	61.4
14 83.9 40.3	14 83.9 40.3	83.9 40.3	40.3	40.3	62	<u></u>	58.2	51.0		45.8	43.6	55.7	65	110	58.2	 	29.8	2.0	0.09
25.4 12 67.9 41.2 46.3 30 80.4 35.0	30 80.4 35.0	80.4 35.0	35.0	35.0	2 2	7 -	48.2	20.5		49.5	48.8	42.2	20 20	100	57.2		31.2	2.1	80.8
48.5 26 87.7 39.1	26 87.7 39.1	87.7 39.1	39.1	39.1	ळी	4	50.6	39.8	49.1	48.8	33.3	50.8	3 8	108	57.5	12.7	32.1 29.5	2.2	63.4 64 9
53.8 10 89.5 47.4	10 89.5 47.4	89.5 47.4	47.4	47.4	l [©]	60.3	59.3	46.7		47.2	45.7	55.5	89	109	54.4	-	34.4	2.2	64.0
53.9 9 82.1 46.9	9 82.1 46.9	82.1 46.9	46.9	46.9	ျ	0.	58.2	51.2	66.4	47.5	38.5	38.6	88	109	56.9	\vdash	30.8	2.1	69.9
36.1	13 /6.2 36.1	7.6.2 36.1	36.1	36.1	ığ l		60.5	48.2		48.2	48.3	54.6	99 i	106	57.7	14.8	30.5	\dashv	68.7
75.7 24.1 29.1	34 777 430	777 420	7.87	7.87	2 0	0 7	4.00	53.3		45.5	39.7	47.4	- 1	108	59.6	\rightarrow	27.5	\dashv	66.2
46.0 31 82.6 41.6	31 82.6 41.6	82.6 41.6	44.0	44.0	1 4	1 4	53.6	55.3	60.0	39.7 73.5	36.9	20.0	20	109	5,7	_	25.9	+	16.8
11-4 49.8 23 59.3 37.8	49.8 23 59.3 37.8	59.3 37.8	37.8	37.8	20	50.9	55.7	49.6		53.6	37.8	46.7	3 8	100	57.5	14 8	30.4 4 %	2 6	2.5
56.4 4 73.2 37.8	4 73.2 37.8	73.2 37.8	37.8	37.8	57	9	60.4	53.2	63.7	53.4	63.5	54.8	8	107	59.7	—	27.2	200	2 0
52.6 16 89.3 40.5	16 89.3 40.5	89.3 40.5	40.5	40.5	48.	1_	56.0	51.7	1	50.6	45.9	46.0	123	107	59.0	—	20.02	27	22.5
53.0 15 93.6 41.6	15 93.6 41.6	93.6 41.6	41.6	41.6	.88	CM.	58.9	54.2	63.4	51.1	34.2	41.1	99	107	58.8	170333	29.7		1 2
22 96.0	22 96.0 34.4	96.0 34.4	34.4	34.4	59.8	-	46.9	54.5		52.3	37.3	41.8	88	109	58.3	33	28.5		70.6
51.2 19 88.7 41.2	19 88.7 41.2	88.7 41.2	41.2	41.2	59	80	47.7	43.6		52.4	45.5	50.8	99	115	58.9	-	27.8	T	64.0
37.1	32 93.8 37.1	93.8 37.1	37.1	37.1	52	6	46.2	37.0	ıı	46.4	34.7	37.5	63	114	57.2	-	26.1	T	20.1
47.4 28 94.7 41.1	28 94.7 41.1	94.7 41.1	41.1	_	ũ	<u></u>	51.1	40.4	58.8	44.2	33.3	38.0	62	114	56.2	15.2	29.2	2.0	9.99
# Locations Mean 80.5 40.1 57.2	40.1	40.1	40.1		57	Ŋ	52.5	49.2	58.1	45.2	41.8	47.3	-	4	6	3	2	2	7
					14.2		13.8	12.8				1							
LSD(.05) - 13.2	. 13.2	- 13.2	- 13.2	13.2	13.2		11.8	10.4	8.8			1							

Table 16. 2002 North Texas Elite (NTE) by locati

Figure 1. Mixograms and other quality data of the TX98D1170 and TX98V9628 compared with standard checks from 2003 TXE tested at three location. The numerical data below each mixogram represent protein (%), test weight (lb bu⁻¹), and kernel weight (mg) in the first row; single kernel hardness index and distribution in the second row.



REPRODUCE LOCALLY. Include form number and edition date on a	Il reproductions. F	ORM APPROVED - OMB No. 0581-005
U.S. DEPARTMENT OF AGRICULTURE AGRICULTURAL MARKETING SERVICE EXHIBIT E STATEMENT OF THE BASIS OF OWNERSHIP	Application is required in order to detect certificate is to be issued (7 U.S.C. 2-confidential until the certificate is issued.	121). The information is held
1. NAME OF APPLICANT(S)	2. TEMPORARY DESIGNATION	3. VARIETY NAME
T. MAINE OF AFFEIDANT(3)	OR EXPERIMENTAL NUMBER	3. VARIETY NAME
Texas Agricultural Experiment Station	TX98V9628	TAM 112
4. ADDRESS (Street and No., or R.F.D. No., City, State, and ZIP, and Country)	5. TELEPHONE (Include area code)	6. FAX (Include area code)
Office of the Director, TAES 2147 TAMU	(979) 845-4747	(979) 458-4765
College Station, TX 77843-2147	7. PVPO NUMBER	
	2006	00274
8. Does the applicant own all rights to the variety? Mark an "X" in the	ne appropriate block. If no, please expla	in. YES NO
		•
9. Is the applicant (individual or company) a U.S. national or a U.S. I	based company? If no, give name of co	ountry. YES NO
10. Is the applicant the original owner? YES	NO If no, please answer one	of the following:
a. If the original rights to variety were owned by individual(s), is YES b. If the original rights to variety were owned by a company(ies) YES	NO If no, give name of count	ry sed company?
11. Additional explanation on ownership (Trace ownership from original contents)	inal breeder to current owner. Use the re	everse for extra space if needed):
The original breeder (that made the cross) is Dr. David Worrall, of the cross. TAES policy and handbook manual provide that all duties are owned by TAES. A copy of this policy is provided for	germplasm and varieties developed by	ilities in Vernon, Texas, at the time its employees in the course of their
PLEASE NOTE:		
Plant variety protection can only be afforded to the owners (not licens	sees) who meet the following criteria:	
If the rights to the variety are owned by the original breeder, that p national of a country which affords similar protection to nationals of a country which affords similar protection.	person must be a U.S. national, national of the U.S. for the same genus and species	of a UPOV member country, or es.
If the rights to the variety are owned by the company which employ nationals of a UPOV member country, or owned by nationals of a genus and species.	yed the original breeder(s), the company country which affords similar protection to	must be U.S. based, owned by onationals of the U.S. for the same
3. If the applicant is an owner who is not the original owner, both the	original owner and the applicant must m	eet one of the above criteria.
The original breeder/owner may be the individual or company who di Act for definitions.	rected the final breeding. See Section 4	1(a)(2) of the Plant Variety Protection
According to the Paperwork Reduction Act of 1995, an agency may not conduct or sponsor, control number. The valid OMB control number for this information collection is 0581-0055, including the time for reviewing the instructions, searching existing data sources, gathering or specific process.	The time required to complete this information collect	tion is estimated to average 0.1 hour per response.

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, gender, religion, age, disability, sexual orientation, marital or family status, political beliefs, parental status, or protected genetic information. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at 202-720-2600 (voice and TDD).

To file a complaint of discrimination, write USDA, Director, Office of Civil Rights, Room 326-W, Whitten Building, 14th and Independence Avenue, SW, Washington, D.C. 20250-9410 or call (202) 720-5964 (voice and TDD). USDA is an equal opportunity provide and employer.

According to the Paperwork Reduction Act of 1995, an agency may not conduct or sponsor, and a person is not required to respond to a collection of information unless it displays a valid OMB control number. The valid OMB control number for this information collection is 0581-0055. The time required to complete this information collection is estimated to average 5 minutes per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information.

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, gender, religion, age, disability, sexual orientation, marital or family status, political beliefs, parental status, or protected genetic information. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at 202-720-2600 (voice and TDD).

To file a complaint of discrimination, write USDA, Director, Office of Civil Rights, Room 326-W, Whilten Building, 14th and Independence Avenue, SW, Washington, DC 20250-9410 or call 202-720-5964 (voice and TDD). USDA is an equal opportunity provider and employer.

U.S. DEPARTMENT OF AGRICULTURE AGRICULTURAL MARKETING SERVICE SCIENCE AND TECHNOLOGY PLANT VARIETY PROTECTION OFFICE BELTSVILLE, MD 20705

EXHIBIT F
DECLARATION REGARDING DEPOSIT

DECLARATION REGARDING DEPOSIT		
NAME OF OWNER (S)	ADDRESS (Street and No. or RD No., City, State, and Zip Code and Country)	TEMPORARY OR EXPERIMENTAL DESIGNATION
Texas Agricultural Experiment Station	Office of the Director, TAES 2147 TAMU College Station, TX 77843-2147	TX98V9628
		VARIETY NAME TAM 112
NAME OF OWNER REPRESENTATIVE (S)	ADDRESS (Street and No. or RD No., City, State, and Zip Code and Country)	FOR OFFICIAL USE ONLY
Mark A. Hussey	Office of the Director, TAES 2147 TAMU College Station, TX 77843-2147	PVPO NUMBER 200600274

I do hereby declare that during the life of the certificate a viable sample of propagating material of the subject variety will be deposited, and replenished as needed periodically, in a public repository in the United States in accordance with the regulations established by the Plant Variety Protection Office.

Mala Dussey

8-10-2006

Date